


# one & done


OTSC® – superior outcome in hemostasis



st  
line



# Results of the 3 RCTs – OTSC® vs. former standard therapy in the first-line treatment of non-variceal UGI bleeding



n [OTSC/Standard]	53 (25/28)	100 (48/52)	190 (93/97)
Primary endpoint	Clinical success defined as no rebleeding ≤ 30 days	Clinical success defined as no rebleeding ≤ 7 days	Clinical success defined as no rebleeding ≤ 30 days
Main inclusion criteria	Ulcer bleeding & Dieulafoy	High risk lesions (cRS ≥ ?)	All NVUGIB
Intervention arm	OTSC arm <sup>‡</sup> vs. Standard arm <sup>‡</sup> (approx. 50 % endoclips / 50 % thermocoag.)	OTSC arm <sup>‡</sup> vs. Standard arm <sup>‡</sup> (approx. 98 % endoclips / 2 % thermocoag.)	OTSC arm <sup>‡</sup> vs. Standard arm <sup>‡</sup> (approx. 60 % thermocoag. / 30 % endoclips / 10 % other combination)

<sup>‡</sup> optional injection therapy allowed

1 Jensen DM, Kovacs T, Ghassemi KA, Kaneshiro M, Gornbein J. Randomized Controlled Trial of Over-the-Scope Clip as Initial Treatment of Severe Nonvariceal Upper Gastrointestinal Bleeding. Clin Gastroenterol Hepatol. 2021 Nov;19(11):2315-2323.e2

2 Meier B, Wannhoff A, Denzer U, Stathopoulos P, Schumacher B, Albers D, Hoffmeister A, Feisthammel J, Walter B, Meining A, Wedi E, Zachaus M, Pickartz T, Küller A, Schmidt A, Caca K. Over-the-scope-clips versus standard treatment in high-risk patients with acute non-variceal upper gastrointestinal bleeding: a randomised controlled trial (STING-2). Gut. 2022 Jul;71(7):1251-1258.

3 Lau J, Tan CH, Sun X, Song H, Li L, Li R, Feng J, Wang B, Leung W.K, Hartley I, Moss AC, Suen BY, Yu Y, Chan FK. Over-the-scope clips (OTSC) versus standard endoscopic treatment in patients with acute non-variceal upper gastrointestinal bleeding. UEG Week Virtual (October 3-5) 2021